

## Author

Heitzmann, Daniel (Heidelberg DE) | Dipl. Ing. (FH)  
Orthopädische Universitätsklinik Heidelberg - Bewegungsanalytik

## Title

Gait pattern differences depending on the level of amputation

## Coauthors

Braatz F., Alimusaj M. , Wolf S. I. Klinik für Orthopädie und Unfallchirurgie, Universitätsklinikum Heidelberg

## Summary

-

## Introduction

-

## Methods

-

## Results

-

## Conclusion

-

## References

- [1] Heitzmann D., M. Günther, Wolf S.I., M. Alimusaj and Braatz F., Strength deficits in trans-tibial amputees, *Gait & Posture* 30 (2009), Supplement 43
- [2] R. Baumgartner, P. Botta, *Amputation und Prothesenversorgung der unteren Extremität*, (1989)
- [3] H. Sadeghi, P. Allard, F. Prince and H. Labelle, Symmetry and limb dominance in able-bodied gait: a review, *Gait Posture* 12 (2000), pp.34-45.
- [4] E. Ludwigs, M. Freslier, M. Bellmann, T. Schmalz and S. Blumentritt (2009) Biomechanische Charakteristiken des Gangbildes verschiedener Amputationsniveaus. Abstractband 6 Jahrestagung der DGfB Münster 71-72.
- [5] M.P. Kadaba, H.K. Ramakrishnan, M.E. Wootten, J. Gainey, G. Gorton and G.V. Cochran, Repeatability of kinematic, kinetic, and electromyographic data in normal adult gait, *J Orthop Res* 7 (1989), 849-860.

[6] R.B. Davis, S. Ounpuu, D. Tyburski and J.R. Gage, A gait analysis data collection and reduction technique, Human Movement Science 10 (1991), 575-587