

**Author**

Sabbagh, Daniel (Lüneburg DE) | Dipl.-Ing.(FH)  
FIOR & GENTZ GmbH - Wissenschaftliche Redaktion

**Title**

The importance of a dorsiflexion stop in KAFOs for patients with weak plantar flexors

**Coauthors**

Fior J, Gentz R

**Summary**

-

**Introduction/Basis**

-

**Scientific Method**

-

**Implementation**

-

**Conclusion**

-

**References**

- [1] Perry J, Mulroy SJ, Renwick SE (1993): The Relationship of Lower Extremity Strength and Gait Parameters in Patients with Post-Polio Syndrome. Archives of Physical Medicine and Rehabilitation 74(2): 165-169
- [2] Lehmann JF, Condon SM, de Lateur BJ, Smith JC (1985): Ankle-Foot Orthoses: Effect on Gait Abnormalities in Tibial Nerve Paralysis. Archives of Physical Medicine and Rehabilitation 66(4): 212-218
- [3] Nollet F, Noppe CT (2008): Orthoses for persons with postpolio syndrome. In: Hsu JD, Michael JW, Fisk JR (Hrsg.): AAOS Atlas of Orthoses and Assistive Devices, 4. Auflage. Philadelphia: Mosby. 411-417
- [4] Ploeger HE, Bus SA, Brehm MA, Nollet F (2014): Ankle-foot orthoses that restrict dorsiflexion improve walking in polio survivors with calf muscle weakness. Gait & Posture 40(3): 391–398

[5] Beekman C, Perry J, Boyd LA, Newsam CJ, Mulroy SJ (2000): The Effects of a Dorsiflexion-Stopped Ankle-Foot Orthosis on Walking in Individuals with Incomplete Spinal Cord Injury.

Topics in Spinal Cord Injury Rehabilitation 5(4): 54–62

[6] Mulroy SJ, Eberly VJ, Gronely JK, Weiss W, Newsam CJ (2010): Effect of AFO Design on Walking after Stroke: Impact of Ankle Plantar Flexion Contracture. Prosthetics and Orthotics

International 34(3): 277-292