

## Author

Schulz, Axel (Frechen DE) | Dr. med.  
Össur Europe B.V. - None

## Title

Recommendation for rehabilitation following cartilage repair - a global cartilage expert consensus

## Coauthors

Albohm M, Carter B, Rembitzki I, Embrechts P, Rogoschin J

## Summary

-

## Introduction

-

## Methods

-

## Results

-

## Conclusion

-

## References

1. Ebert et al. (2017) Two-Year Outcomes of a Randomized Trial Investigating a 6-Week Return to Full Weightbearing After Matrix-Induced Autologous Chondrocyte Implantation.[HYPERLINK I](#) "Am J Sports Med. 2017 Mar;45(4):838-848.
2. Marder RA, Hopkins GJ, Timmerman LA. Arthroscopic microfracture of chondral defects of the knee: a comparison of two postoperative treatments. Arthroscopy. 2005;21:152–158.
3. Steadman JR, Rodkey WG, Briggs KK. Microfracture to treat full-thickness chondral defects: surgical technique, rehabilitation, and outcomes. J Knee Surg. 2002;15: 170–176.
4. Wondrasch et al. (2015) Am J Sports Med. 2015 Jan;43(1):146-53. Effect of accelerated weightbearing after matrix-associated autologous chondrocyte implantation on the femoral condyle: a prospective, randomized controlled study presenting MRI based and clinical outcomes after 5 years